Astronomy & Astrophysics manuscript no. main
June 4, 2010

© ESO 2010

arXiv:1006.0588v1 [astro-ph.SR] 3 Jun 2010

A survey of T Tauri stars with AKARI toward the Taurus-Auriga region

Satoshi Takita[1,2], Hirokazu Kataza[2], Yoshimi Kitamura[2], Daisuke Ishihara[3], Yoshifusa Ita[4] ⋆, Shinki Oyabu[2] ⋆⋆, and Munetaka Ueno[2]

[1] Department of Space and Astronautical Science, The Graduate University for Advanced Studies (Sokendai), 3-1-1 Yoshinodai, Chuo, Sagamihara, Kanagawa 229-8510, Japan
[2] Institute of Space and Astronautical Science, Japan Aerospace Exploration Agency, 3-1-1 Yoshinodai, Chuo, Sagamihara, Kanagawa 229-8510, Japan
[3] Division of Particle and Astrophysical Sciences, Nagoya University, Furo-cho, Chikusa-ku, Nagoya, 464-8602, Japan
[4] National Astronomical Observatory of Japan, 2-21-1 Osawa, Mitaka, Tokyo 181-8588, Japan

Received 2 November 1992 / Accepted 7 January 1993

ABSTRACT

Context. AKARI, the first Japanese infrared astronomical satellite, has completed an All-Sky Survey at the mid- to far-infrared wavelengths with greater spatial resolutions and sensitivities than those of the previous survey with Infrared Astronomical Satellite (IRAS).
Aims. We search new T Tauri star (TTS) candidates with the mid-infrared (MIR) part of the AKARI All-Sky Survey at 9 and 18 μm wavelengths.
Methods. We used the point source catalogue (PSC), obtained by the Infrared Camera (IRC) on board AKARI. We combined the 2MASS PSC and the 3rd version of the USNO CCD Astrograph Catalogue (UCAC) with the AKARI IRC-PSC, and surveyed 517 known TTSs over a 1800-square-degree part of the Taurus-Auriga region to find criteria to extract TTSs. We considered asymptotic giant branch (AGB) stars, post-AGB stars, Planetary Nebulae (PNe), and galaxies, which have similar MIR colours, to separate TTSs from these sources. We finally searched new TTS candidates from AKARI IRC-PSC in the same Taurus-Auriga region.
Results. Of the 517 known TTSs, we detected 133 sources with AKARI: 46 sources were not detected by IRAS. Based on the colour-colour and colour-magnitude diagrams made from the AKARI, 2MASS, and UCAC surveys, we propose the criteria to extract TTS candidates from the AKARI All-Sky data, and 68/133 AKARI detected TTSs have passed these criteria. On the basis of our criteria, we selected 176/14725 AKARI sources as TTS candidates which are located around the Taurus-Auriga region. Comparing these sources with SIMBAD, there are 148 previously identified sources including 115 Young Stellar Objects (YSOs), and 28 unidentified sources.
Conclusions. Based on SIMBAD identifications, we take the TTS-identification probability using our criteria to be ~75 %. We find 28 TTS candidates, of which we expect ~21 to be confirmed once follow-up observations can be obtained. Although the probability of ~75 % is not so high, it is affected by the completeness of the SIMBAD database, and we can search for TTSs over the whole sky, over all star forming regions.

Key words. Stars: formation — Stars: late-type — Stars: pre-main sequence — infrared: stars

1. INTRODUCTION

T Tauri stars (TTSs) are low-mass pre-main sequence (PMS) stars with ages of ~ 10^6–10^7 yrs (Adams et al. 1988). TTSs were traditionally identified with their strong Hα emission line toward their natal molecular clouds, i.e., classical TTSs (CTTSs) with ages of ~ 10^6 yr. On the other hand, X-ray observations, such as the Einstein Observatory and ROSAT surveys, have discovered another class of PMS stars (e.g., Neuhäuser et al. 1995a). Since these sources have the "weak" Hα emission line with equivalent widths of ≤ 10 Å, they are called weak-line TTSs (WTTSs) and have ages of ~ 10^7 yr. The youth of WTTSs was confirmed by strong Li ɪ absorption lines, because lithium is easily destroyed in the stellar atmosphere with high temperature. In contrast to the CTTS case, many WTTSs are found outside the molecular clouds. This situation is interpreted as follows: The natal molecular clouds have been already dispersed at ~ 10^7 yr, or the stars have left the natal clouds owing to their motions with a few mas/yr.

The circumstellar disks around TTSs are believed to be the birthplaces of planets. The pioneering studies of the disks are the IRAS and millimetre continuum surveys towards the Taurus-Auriga region (Strom et al. 1989; Beckwith et al. 1990). Strom et al. (1989) found that about a half of CTTSs have excess emission at the infrared (IR) wavelength stronger than that expected from their photospheres. The excess emission is well interpreted as the thermal emission from the circumstellar disks heated by the central stars and/or mass accretion. On the other hand, most WTTSs lack such excess emission. Even the recent IR observations with the *Spitzer* Space Telescope have confirmed such a low detection rate of the excess emission towards WTTSs. Silverstone et al. (2006) showed that 5 of 74 young stars (3–30 Myr) have strong IR excess, but they have spectral

---

[⋆] Present address is Astronomical Institute, Graduate School of Science, Tohoku University, 6-3 Aramaki, Aoba-ku, Sendai, 980-8578, Japan

[⋆⋆] Present address is Subaru Telescope, National Astronomical Observatory of Japan, 650 North A'ohoku Place, Hilo, HI 96720, USA energy distributions consistent with CTTSs. Padgett et al. (2006) observed 83 WTTSs outside their natal clouds, and found only 5 WTTSs have excess emission. The largest *Spitzer* survey of WTTSs so far was done by Cieza et al. (2007), which observed more than 230 WTTSs located in the Ophiuchus, Lupus, and Perseus molecular clouds. Their data indicate that ∼20 % of the WTTSs have IR excess emission, but no IR excess for the stars older than 10 Myr. Consequently, it is most likely that the disk dissipation time scale is about 10 Myr. However, IRAS did not have enough sensitivities to detect the excess emission of WTTSs, and *Spitzer* cannot cover "all" WTTSs. Therefore, we need unbiased and high-sensitivity surveys to study the WTTS disks with good statistics.

2. AKARI all-sky data

2.1. AKARI IRC All-Sky Survey

AKARI is the first Japanese infrared astronomical satellite dedicated to infrared astronomy (Murakami et al. 2007). One of the major observation programs of AKARI is an All-Sky Survey at the mid- to far-infrared wavelengths with 6 photometric bands. AKARI has a higher sensitivity, a higher spatial resolution, and a wider wavelength coverage than those of the previous IRAS survey. The mid-infrared (MIR) survey has been carried out with the *S9W* (9 μm) and *L18W* (18 μm) bands using the Infrared Camera (IRC; Onaka et al. 2007). The 5 σ detection limit for a point source is estimated to be 50 and 120 mJy at the *S9W* and *L18W* bands, respectively. The spatial resolution is around 5′′. More than 96 % of the entire sky has been observed with the two bands. The first version of the AKARI IRC point source catalogue (hereafter IRC-PSC) was publicly released in March 2010. The analysis of this paper is based on the 1st version of the IRC-PSC. The details of the AKARI IRC All-Sky Survey and its data reduction processes are described in Ishihara et al. (2010).

2.2. Comparison with the NIR 2MASS and optical UCAC catalogues

We compared the IRC-PSC with the 2MASS PSC by a near-infrared (NIR) survey (Skrutskie et al. 2006) using a simple positional correlation method. We used a positional tolerance of 5′′, which is the spatial resolution of the AKARI IRC All-Sky Survey. More than 99 % of the AKARI sources agree well with those in the 2MASS PSC within the accuracy. We also compared the IRC-PSC with the 3rd version of the USNO CCD Astrograph Catalogue (UCAC) by an optical survey (Zacharias et al. 2010) in the same way. About 70 % of the AKARI sources have optical counterparts within the 5′′ accuracy. This relatively low cross-identification rate comes from the limited magnitude range in the UCAC survey: Most nearby (< 100 pc) stars are saturated, and distant (> 1 kpc) or heavily reddened stars have no UCAC entry because of its sensitivity. Since the UCAC catalogue contains "stars" with the magnitude range of R = 7.5–16.3 in a 579–642 nm band, it is useful to search stars at distances of about 100 pc, which is the typical distance to nearby molecular clouds (for example, if we put the Sun at 140 pc, the magnitude becomes ∼11).

3. AKARI IRC observations of the previously known T Tauri stars in the Taurus-Auriga region

3.1. Previously known members in the Taurus-Auriga region

We selected a 1800-square-degree ($2^{\mathrm{h}}40^{\mathrm{m}} < \mathrm{R.A.} < 5^{\mathrm{h}}40^{\mathrm{m}}$ and $0^\circ < \mathrm{Dec} < 40^\circ$) part of the Taurus-Auriga region for this study. The Taurus-Auriga region is a well-studied low-mass star-forming one at a close distance of ∼ 140 pc with hundreds of pre-main-sequence (PMS) stars. We first compiled known Taurus PMS members from Rebull et al. (2010), Kenyon et al. (2008), Güdel et al. (2007), Beckwith et al. (1990), and Strom et al. (1989), which are associated with the Taurus molecular cloud. Second, we picked up TTSs from the PMS objects as follows: (1) Class II/III objects including the 'new', 'probable', and 'possible' members in Rebull et al. (2010); (2) Objects listed in Kenyon et al. (2008) that were confirmed as Class II/III objects on the basis of the SIMBAD database and their references; (3) Objects which are labelled as CTTSs or WTTSs in Güdel et al. (2007). Note that Beckwith et al. (1990) and Strom et al. (1989) listed only TTSs. Furthermore, we added TTSs which are discovered by the Roentgen satellite (ROSAT) and distributed from the cloud (Li & Hu 1998; Magazzù et al. 1997; Wichmann et al. 1996). In our TTS list, we considered close (< 5′′; the spatial resolution of the AKARI IRC All-Sky Survey) binaries as a single source. Finally, we got a catalogue of 517 TTSs in the selected region. The distribution of our input TTSs is shown in Figure 1.

3.2. Statistical Properties of the previously known T Tauri stars detected with AKARI

We totally detected 133 TTSs with AKARI within the positional accuracy of 5′′, i.e., the resolution of the AKARI IRC All-Sky Survey: 127 and 95 sources were detected at the *S9W* and *L18W* bands, respectively. There are 6 sources which were detected only with the *L18W* band: 3 sources (IRAS 04200+2759, IRAS 04295+2251, and UX Tau) were observed/detected only once at the *S9W* band, so the IRC-PSC does not contain these sources, and the other 3 sources were not detected at all. The positions of almost all the AKARI detected sources agree well with those in the previous catalogues within the uncertainty of 3′′ (see Figure 2). Although GI Tau has larger separations than 3′′, it still satisfy the positional accuracy of AKARI IRC-PSC. The photometric data of the AKARI detected TTSs are listed in Table 1.

In the 133 TTSs, 46 faint sources have no counterparts in the IRAS PSC (FQUAL12 = 3 or FQUAL25 = 3) with a searching radius of 60′′, the IRAS resolution, as shown in Figure 3. These 'new detections' are due to the higher sensitivity and spatial resolution of AKARI than those of IRAS. Actually, Figure 3 shows that the detection limit for the previously known Taurus TTSs at the *S9W* band has been improved: IRAS could detect almost all the TTSs that have brighter *S9W* and *L18W* band magnitudes than 6 and 4, respectively, but only about a quarter of the fainter sources were detected with IRAS. Figure 4 shows the $(K_S - S9W)$ v.s. $(S9W)$ colour-magnitude diagram of the AKARI detected sources. Since about the half of the sources that were not detected with IRAS have small colours with $K_S - S9W < 1$ and most of the IRAS detected TTSs have larger colours with $K_S - S9W > 2$, there seems to exist a gap at $K_S - S9W \sim 0.5$. The sources with $K_S - S9W < 1$ are most likely to be WTTSs, because they have weak Hα emission (except HT Tau, whose Hα equivalent width is not given) and are located near the periphery of the clouds or outside the clouds. AKARI has succeeded in de-



Fig. 1. Distribution of the target TTSs (small open circles). The small filled circles indicate the TTSs detected with AKARI and the large open ones mean the IRAS detection, on the IRAS 100 μm map.

tecting WTTSs with weak excess emission at the sensitive *S9W* band.

There are 18 TTSs which are not catalogued in the IRC-PSC within a 5″ searching radius, but are catalogued in the IRAS PSC within 60″ radius. Of these sources, 15 sources have AKARI counterparts within the positional accuracy of the IRAS, but the remaining 3 sources have no counterparts in the IRC-PSC. IRAS 04302+2247 was observed once or four times at the *S9W* or *L18W* bands, respectively, but not detected. Although this source is a well-known TTS with an edge-on disk, it is faint at the *S9W* and *L18W* bands, suggesting the presence of an inner gap in the disk.. Indeed, it has bright 24 μm magnitude of 3.57 but faint 8 μm one of 9.71 (Rebull et al. 2010). IRAS 04216+2603 was observed four times and detected only once at the *L18W* band, and not observed at the *S9W* band at all. Since the IRC-PSC requires more than one detection for a 'real' point source to reject a moving object or a fake one, the source is not catalogued in the IRC-PSC. Finally, DM Tau was detected twice at the *L18W* band, but the positions in the two independent images have a larger difference than 5″ between each other under the current positional accuracy, i.e., no entry in the IRC-PSC.

Although Spitzer didn't cover the entire region we searched, sources inside the Taurus molecular cloud and some other sources outside the cloud have been also observed with Spitzer



Fig. 2. Histogram of the positional differences between the TTSs detected with AKARI in the IRC-PSC and those in the previous catalogues. The size of the positional difference bin is 0.1 arcsec.



Fig. 3. Histogram of the *S9W* (*top*) and *L18W* (*bottom*) magnitudes of the TTSs detected with AKARI. The filled and open bars indicate the sources that were detected and not detected with IRAS, respectively. The size of the magnitude bin is 0.25 magnitude.

(e.g., Rebull et al. 2010). Since Spitzer has higher sensitivity than AKARI All-Sky Survey, Spitzer has discovered more faint sources as shown in Figure 5, which shows the histogram of the IRAC4 (8 μm) magnitudes of the detected TTSs with Spitzer. The AKARI All-Sky Survey could detect more than ~90% of the sources with magnitude brighter than 7.5 at the IRAC4 band, which agrees well with the detection limit of the AKARI *S9W* band.

4. How to extract TTS candidates from the AKARI All-Sky data

4.1. Other types of sources in the whole sky

Since the following types of sources are known to have similar colours to those of the TTSs, we should reveal the colour properties of the sources. We considered the four additional catalogues of (1) asymptotic giant branch (AGB) stars of 126 carbon and 563 OH/IR stars (Le Bertre et al. 2003), (2) 326 Post-AGB stars (Szczerba et al. 2007), (3) 1143 Planetary Nebulae (PNe) (Acker et al. 1994), and (4) 2907 extragalactic objects which have brighter flux density than 100 mJy at the IRAS 12 μm band (the NASA Extragalactic Database). In contrast to the TTS case, we examined these sources in the whole sky because of a small number of the sources toward the Taurus-Auriga region.



Fig. 4. $(K_S - S9W)$ v.s. $(S9W)$ colour-magnitude diagram of the AKARI detected PMS stars. The filled and open circles indicate the sources that were detected and not detected with IRAS, respectively. The arrow shows the interstellar extinction vector of $A_V = 20$ mag, using the Weingartner & Draine (2001) Milky Way model of $R_V = 3.1$.



Fig. 5. Histogram of the *IRAC4* magnitudes of the PMS stars which are listed in Table 4 by Rebull et al. (2010). The filled and open bars indicate the sources that were detected and not detected with AKARI, respectively. The size of the magnitude bin is 0.25 magnitude.

4.2. Colour-colour diagrams

Figure 6 shows the $(K_S - S9W)$ v.s. $(J - K_S)$ and $(K_S - L18W)$ v.s. $(J - K_S)$ colour-colour diagrams, where the *S9W* and *L18W* excess emission can be clearly recognized. One object, 1RXS J032409.7+123745, does not seem to have excess emission at both the *S9W* and *L18W* bands among the TTSs in the Taurus-Auriga region. Although this source is listed as a WTTS with spectral type of K2 (Li & Hu 1998), it seems to be a Li-rich giant star because of high luminosity ($J \sim 4$ and $V \sim 6$) not expected for a K2 dwarf star at a distance of ~ 140 pc. On the other hand, the other types of the sources, which are stars surrounded with dust, also have significant IR excess emission. Therefore, we need to separate these sources from TTSs in the IRC-PSC.

First, we separate almost all carbon and OH/IR stars from the TTSs in the $(S9W - L18W)$ v.s. $(J - K_S)$ colour-colour diagram shown in Figure 7. Since an AGB star typically has hot dust near its photosphere, the $(S9W - L18W)$ colour, which represents the dust temperature, becomes *blue*. On the other hand, since the majority of the dust around a TTS is cold, the MIR colour is *red*. To remove about 80 and 70 % of carbon and OH/IR stars, we propose the first criterion as

$$J - K_S \leq 7.7 \times (S9W - L18W) - 8.5. \quad (1)$$



Fig. 6. *Top*: ($K_S - S9W$) v.s. ($J - K_S$) colour-colour diagram. The grey dots with error bars indicate all the AKARI point sources in the selected Taurus-Auriga region. The red plus and cross symbols show carbon and OH/IR stars in the whole sky, respectively. The magenta diamonds, the orange circles, and the blue squares are post-AGB stars, PNe, and extragalactic objects in the whole sky, respectively. The green stars mean the TTSs in the Taurus-Auriga region. The arrow shows the interstellar extinction vector of A_V = 20 mag, estimated from the Weingartner & Draine (2001) Milky Way model of R_V = 3.1. *Bottom*: ($K_S - L18W$) v.s. ($J - K_S$) colour-colour diagram. The symbols and arrow are the same as in the *top* panel.

However, post-AGB stars and PNe have the *red* MIR colour because they have cold dust. Furthermore, galaxies also have cold dust. Therefore, we can separate only 15 % of post-AGB stars, and few PNe and galaxies from TTSs. Note that this criterion is valid for the sources that were detected at both the *S9W* and *L18W* bands. We do not remove the sources which were detected at only one band.

4.3. Colour-magnitude diagram

Second, we can separate post-AGB stars, PNe, and galaxies from TTSs in the ($UCAC - J$) v.s. ($UCAC$) colour-magnitude diagram. Owing to the low luminosities of PNe (white dwarfs) and galaxies at the visible wavelengths, we can remove about 90 and 97 % of PNe and galaxies, respectively, by the following 2nd criterion:

$$UCAC \leq 1.5 \times (UCAC - J) + 10. \qquad (2)$$



Fig. 7. ($S9W - L18W$) v.s. ($J - K_S$) colour-colour diagram. The symbols and arrow are the same as in Figure 6. The black solid line indicates our criterion (1). The black broken lines indicate $S9W - L18W$ colours of 200, 500, and 1000 K blackbody.



Fig. 8. ($UCAC - J$) v.s. ($UCAC$) colour-magnitude diagram. The symbols are the same as in Figure 6. The arrow shows the interstellar extinction vector of A_V = 5 mag, estimated from the Weingartner & Draine (2001) Milky Way model of R_V = 3.1. The black line indicates our criterion (2).

From this criterion, we can also get rid of 26 % of post-AGB stars. Furthermore, 4 and 12 % of carbon and OH/IR stars, respectively, can be also separated. Considering the detection limits of the IRC-PSC and the UCAC, this method seems effective only for extracting nearby ($\lesssim$ 100 pc) young sources. Note that we never pick up TTSs with their edge-on disks, because they are not optically visible.

Figure 9 shows the same colour-colour diagrams as in Figure 6, but for the remaining sources that could not be removed by the criteria (1) and (2); most of the remaining sources with the excess emission are the TTSs. Finally, we propose the following two criteria to pick up most of the TTSs with less contamination of other types of sources:

$$-11.5 \leq (J - K_S) - 2.5 \times (K_S - S9W) \leq -2.5 \qquad (3)$$
$$\&\& \; 0.5 \leq J - K_S \leq 3.5,$$
$$-12 \leq (J - K_S) - 1.9 \times (K_S - S9W) \leq -5 \qquad (4)$$
$$\&\& \; 0.5 \leq J - K_S \leq 3.5.$$

Note that we cannot pick up TTSs which have weak MIR excess emission, because they are contaminated by field stars. Consequently, most of AGB stars, PNe, and galaxies have been removed with our criteria. However, 34 % of post-AGB stars still remain, and we need follow-up observations to distinguish young stars from post-AGB stars. Of the 133 AKARI detected TTSs, 68 sources have passed these criteria.



Fig. 9. Colour-Colour diagrams, the same as in Figure 6, but for the selected sources with the criteria (1) and (2) in the text. The green parallelograms indicates our criteria (3) and (4).

5. Evaluation of the criteria

5.1. New TTS candidates from our criteria

On the basis of our criteria, we selected 176 sources out of 14725 AKARI sources, which are distributed in the area between 2^h40^m and 5^h40^m in Right Ascension and between 0° and 40° in Declination. We farther searched these sources in SIMBAD with a $10''$ searching radius. In the 176 selected sources, 21 sources could not be found in the SIMBAD database: there were 115 YSOs, 4 Be stars, 10 AGB stars, 1 Galaxy, 18 other types of objects (mostly variable stars), and the remaining sources consist of 7 unknown objects. Since ~75 % of the SIMBAD identified sources were YSOs, we take the TTS-identification probability based on our criteria to be 75 %, though that is of course affected by the completeness of the SIMBAD database.

We now have found the 21 new TTS candidates in addition to the 7 unknown sources. The infrared properties of the 28 TTS candidates are listed in Table 2. Considering the TTS-identification probability of 75 %, we expect 21 new TTSs. We need follow-up observations to distinguish whether these sources are really TTSs or not.

5.2. Comparison with other methods

Many astronomers have discussed about the criteria to extract YSO candidates from their surveyed data. Recently, Evans et al. (2009) and Rebull et al. (2010) surveyed YSOs in nearby clouds using Spitzer. Evans et al. (2009) surveyed 5 clouds (15.5 deg^2 region in total) and listed ~1000 YSO candidates based on the colour-magnitude and colour-colour diagrams by 2MASS and Spitzer. They showed that there could be 51 galaxies in their YSO candidates. However, since their surveyed areas are highly embedded ($A_V > 2$) and are located at high galactic latitudes, they paid less attention to AGB stars. Rebull et al. (2010) surveyed a ~44 deg^2 region of the Taurus molecular cloud, and listed 148 new candidate Taurus members based on colours and magnitudes by 2MASS and Spitzer together with the images of SDSS and CHFT. Of these 148 new candidates, they did follow-up spectroscopic observations for about half of the sources, and confirmed 47 new Taurus candidates, 7 extragalactic objects, and 1 Be star; the remaining 93 sources await further follow-up observations.

These two studies have the higher (973/1024, Evans et al. 2009; 47/55, Rebull et al. 2010) YSO-identification probabilities than our probability of ~75 %. This is mainly because we have data at only 2 bands; they usually have more than 4 bands. Furthermore, because the detection limits of the AKARI All-Sky Survey are relatively shallower than those of the pointed observations of Spitzer, it is difficult to detect MIR faint objects. However, since AKARI observed almost the whole sky, we can find TTS candidates toward all nearby star forming region.

Acknowledgements. This work is based on observations with AKARI, a JAXA project with the participants of ESA. We gratefully acknowledge all the members of the AKARI project for their support on this project. This publication makes use of data products from the Two Micron All Sky Survey, which is a joint project of the University of Massachusetts and the Infrared Processing and Analysis Center/California Institute of Technology, funded by the National Aeronautics and Space Administration and the National Science Foundation. This research has made use of the SIMBAD database and the VizieR catalogue access tool, CDS, Strasbourg, France This research has made use of the NASA/IPAC Extragalactic Database (NED) which is operated by the Jet Propulsion Laboratory, California Institute of Technology, under contract with the National Aeronautics and Space Administration.

References

Acker, A., Ochsenbein, F., Stenholm, B., et al., 1994, VizieR Online Data Catalog, 5084, 0
Adams, F. C., Lada, C. J., & Shu, F. H., 1988, ApJ, 326, 865
Beckwith, S. V. W., Sargent, A. I., Chini, R. S., & Guesten, R., 1990, AJ, 99, 924
Beichman, C. A., Neugebauer, G., Habing, H. J., Clegg, P. E., & Chester, T. J., 1988, Infrared Astronomical Satellite (IRAS) Catalogs and Atlases, NASA RP-1190
Carpenter, J. M., Bouwman, J., Mamajek, E. E., et al., 2009, ApJS, 181, 197
Cieza, L., Padgett, D. L., Stapelfeldt, K. R., et al. 2007, ApJ, 667, 308
Cohen, M. & Kuhi, L. V., 1979, ApJS, 41, 743
Evans, N. J., Dunham, M. M., Jørgensen, J. K., et al., 2009, ApJS, 181, 321
Furlan, E., Hartmann, L., Calvet, N., et al., 2006, ApJS, 165, 568
Güdel, M., Briggs, K. R., Arzner, K., et al., 2007, A&A, 468, 353
Harris, S., Clegg, P., & Hughes, J., 1988, MNRAS, 235, 441
Harvey, P., Merín, B., Huard, T. L., et al., 2007, ApJ, 663, 1149
Hayashi, C., Nakazawa, K., & Nakagawa, Y., 1985, Protostars and Planets II, 1100
Herbig, G. H. & Bell, K. R., 1988, Third Catalog of Emission-Line Stars of the Orion Population, Lick Observatory Bulletin, No. 1111

Ishihara, D., Onaka, T., Kaneda, H., et al., 2007, PASJ, 59, S443
Ishihara, D., Onaka, T., Kataza, H., et al., 2010, A&A, 514, A1
Ita, Y., Matsuura, M., Ishihara, D., et al., 2010, A&A, 514, A2
Kenyon, S. J., Gómez, M., & Whitney, B. A., Handbook of Star Forming Regions, Volume I, 405
Kitamura, Y., Momose, M., Yokogawa, S., et al., 2002, ApJ, 581, 357
Le Bertre, T., Tanaka, M., Yamamura, I., & Murakami, H., 2003, A&A, 403, 943
Li, J. Z. & Hu, J. Y., 1998, A&AS, 132, 173
Luhman, K. L., Whitney, B. A., Meade, M. R., et al., 2006, ApJ, 647, 1180
Luhman, K. L., Mamajek, E. E., Allen, P. R., & Cruz, K. L., 2009, ApJ, 703, 399
Luhman, K. L., Allen, P. R., Espaillat, C., Hartmann, L., & Calvat, N., 2010, ApJS, 186, 111
Magazzù, A., Martín, E. L., Sterzik, M. F., et al., 1997, A&AS, 124, 449
Murakami, H., Baba, H., Barthel, P., et al., 2007, PASJ, 59, S369
Neuhäuser, R., Sterzik, M. F., Schmitt, J. H. M. M., Wichmann, R., & Krautter, J., 1995a, A&A, 295, L5
Neuhäuser, R., Sterzik, M. F., Schmitt, J. H. M. M., Wichmann, R., & Krautter, J., 1995b, A&A, 297, 391
Neuhäuser, R., Sterzik, M. F., Schmitt, J. H. M. M., Wichmann, R., & Krautter, J., 1995c, A&A, 299, L13
Onaka, T., Yamamura, I., de Jong, T., et al., 1998, Ap&SS, 255, 331
Onaka, T., Matsuhara, H., Wada, T., et al., 2007, PASJ, 59, S401
Padgett, D. L., Cieza, L., Stapelfeldt, K. R., et al., 2006, ApJ, 645, 1283
Rebull, L. M., Padgett, D. L., McCabe, C. -E., et al., 2010, ApJS, 186, 259
Silverstone, M. D., Meyer, M. R., Mamajek, E. E., et al., 2006, ApJ, 639, 1138
Skrutskie, M. F., Cutri, R. M., Stiening, R., et al., 2006, ApJ, 131, 1163
Strom, K. M., Strom, S. E., Edwards, S., Cabrit, S., & Skrutskie, M. F. 1989, AJ, 97, 1451
Szczerba, R., Siódmiak, N., Stasińska, G., et al., 2007, A&A, 469, 799
Tanabé, T., Sakon, I., Cohen, M., et al., 2008, PASJ, 60, S375
Weingartner, J. C. & Draine, B. T., 2001, ApJ, 548, 296
Wichmann, R., Krautter, J., Schmitt, J. H. M. M., et al. 1996, A&A, 312, 439
Wyatt, M. C., Dent, W. R. F., & Greaves, J. S., 2003, MNRAS, 342, 876
Zacharias, N., Urban, S. E., Zacharias, M. I., et al., 2004, AJ, 127, 3043
Zacharias, N., Finch, C., Girard, T., et al., 2010, AJ, 139, 2184

Table 1. AKARI, UCAC, and 2MASS photometric data for the previously known Taurus members.

AKARI IRC name[a]	Identifier	AKARI *S9W*[b] [mag]	AKARI *L18W*[b] [mag]	UCAC[c] [mag]	2MASS *J* [mag]	2MASS *H* [mag]	2MASS K_S [mag]
0311138+222457	1RXS J031113.0+222518	6.847 ± 0.154		8.359 ± 0.093	7.273 ± 0.020	7.069 ± 0.053	6.968 ± 0.027
0324065+234706	1RXS J032406.6+234714	6.008 ± 0.048		10.018 ± 1.000	7.128 ± 0.020	6.492 ± 0.020	6.267 ± 0.018
0324102+123746	1RXS J032409.7+123745[d]	2.950 ± 0.017	2.902 ± 0.023	6.841 ± 0.484	4.097 ± 0.292	3.490 ± 0.248	3.394 ± 0.244
0327325+255400	1RXS J032733.2+255405	6.947 ± 0.508		9.090 ± 1.000	7.927 ± 0.032	7.550 ± 0.051	7.488 ± 0.023
0400095+081818	RXJ0400.1+0818N	7.041 ± 0.183		9.775 ± 0.048	8.281 ± 0.024	8.028 ± 0.000	7.843 ± 0.000
0404430+261856	L1489 IRS	3.745 ± 0.020	0.694 ± 0.022		14.129 ± 0.000	11.908 ± 0.000	9.859 ± 0.043
0406513+254127	RXJ0406.8+2541	6.207 ± 0.126		11.452 ± 0.058	8.767 ± 0.021	8.026 ± 0.034	7.750 ± 0.018
0413271+281624	Anon 1	6.785 ± 0.221		13.261 ± 0.171	8.826 ± 0.030	7.794 ± 0.027	7.460 ± 0.029
0413532+281124	IRAS 04108+2803 A	6.022 ± 0.116			13.639 ± 0.024	11.521 ± 0.024	10.370 ± 0.019
0413573+291820	IRAS 04108+2910	5.648 ± 0.044	3.559 ± 0.152	16.015 ± 0.316	11.310 ± 0.023	10.157 ± 0.024	9.356 ± 0.024
0414129+281212	V773 Tau ABC	4.213 ± 0.028	2.247 ± 0.039	10.434 ± 0.033	7.494 ± 0.024	6.644 ± 0.020	6.210 ± 0.047
0414135+281249	FM Tau	5.729 ± 0.057	3.554 ± 0.036		10.329 ± 0.021	9.391 ± 0.022	8.763 ± 0.022
0414146+282758	FN Tau	5.383 ± 0.096	2.780 ± 0.039	14.151 ± 0.267	9.469 ± 0.022	8.669 ± 0.036	8.189 ± 0.018
0414170+281058	CW Tau	4.144 ± 0.043	2.324 ± 0.049	13.390 ± 0.169	9.557 ± 0.022	8.243 ± 0.027	7.127 ± 0.023
0414176+280609	CIDA-1	6.126 ± 0.126	4.187 ± 0.156		11.726 ± 0.021	10.584 ± 0.022	9.877 ± 0.021
0414263+280602	IRAS 04113+2758 A	3.525 ± 0.017	1.031 ± 0.067		12.475 ± 0.029	9.878 ± 0.033	7.777 ± 0.031
0414305+280514	MHO 3	4.049 ± 0.028	1.185 ± 0.022		11.178 ± 0.021	9.249 ± 0.022	8.243 ± 0.023
0414478+264810	CX Tau	6.227 ± 0.093	3.725 ± 0.100	13.589 ± 0.073	9.867 ± 0.022	9.054 ± 0.024	8.807 ± 0.019
0414479+275234	LkCa 3 AB	6.860 ± 0.189		11.865 ± 0.063	8.363 ± 0.032	7.625 ± 0.023	7.423 ± 0.021
0414492+281230	FO Tau AB	5.719 ± 0.105	3.511 ± 0.052	15.099 ± 0.192	9.650 ± 0.021	8.568 ± 0.021	8.121 ± 0.031
0417337+282047	CY Tau	6.270 ± 0.081		13.022 ± 0.069	9.828 ± 0.020	8.970 ± 0.024	8.597 ± 0.021
0417496+282936	V410 X-ray 1	5.997 ± 0.086	4.415 ± 0.049	16.680 ± 0.515	11.020 ± 0.021	9.733 ± 0.027	9.081 ± 0.019
0418107+251957	[GBM90] L1506 1	6.537 ± 0.096		13.039 ± 0.223	10.744 ± 0.024	9.589 ± 0.022	9.032 ± 0.019
0418310+282716	V410 Tau ABC	7.184 ± 0.111		10.366 ± 0.003	8.449 ± 0.018	7.789 ± 0.021	7.629 ± 0.026
0418311+281629	DD Tau AB	4.271 ± 0.027	2.366 ± 0.043	14.748 ± 0.232	9.830 ± 0.021	8.682 ± 0.020	7.878 ± 0.020
0418316+281658	CZ Tau AB	4.471 ± 0.033	2.461 ± 0.046	15.239 ± 0.023	10.517 ± 0.021	9.774 ± 0.022	9.362 ± 0.017
0418320+283115	IRAS 04154+2823	4.882 ± 0.102	2.420 ± 0.048		15.187 ± 0.048	12.369 ± 0.022	10.274 ± 0.017
0418344+283029	V410 X-ray 2		4.559 ± 0.071		13.773 ± 0.026	10.708 ± 0.022	9.215 ± 0.017
0418406+281914	V892 Tau	1.278 ± 0.016	-1.777 ± 0.016	14.846 ± 0.111	8.742 ± 0.027	7.016 ± 0.029	5.787 ± 0.016
0418470+282008	Hubble 4	6.931 ± 0.275		12.181 ± 0.093	8.560 ± 0.020	7.636 ± 0.026	7.293 ± 0.024
0418514+282026	CoKu Tau/1	5.239 ± 0.089	1.884 ± 0.022		12.866 ± 0.030	11.489 ± 0.031	10.974 ± 0.024
0419010+281942	V410 X-ray 6		4.747 ± 0.066		10.527 ± 0.022	9.597 ± 0.024	9.129 ± 0.021
0419128+282933	FQ Tau AB	7.175 ± 0.294		15.097 ± 0.195	10.491 ± 0.026	9.702 ± 0.024	9.313 ± 0.022
0419158+290626	BP Tau	5.335 ± 0.123	3.303 ± 0.049	11.927 ± 0.081	9.098 ± 0.037	8.220 ± 0.024	7.736 ± 0.023
0419354+282721	FR Tau	6.853 ± 0.357		16.321 ± 0.900	10.954 ± 0.021	10.374 ± 0.022	9.971 ± 0.018
0420258+281922	SST Tau 042025.8+281923	6.564 ± 0.132	4.612 ± 0.054		15.327 ± 0.039	13.422 ± 0.026	11.718 ± 0.021
0420260+280409	SST Tau 042026.0+280408	6.643 ± 0.242	3.740 ± 0.071	14.696 ± 0.190	10.612 ± 0.021	9.949 ± 0.022	9.697 ± 0.018
0421079+270220	CFHT-19	5.309 ± 0.030	2.735 ± 0.054		13.855 ± 0.024	12.062 ± 0.026	10.543 ± 0.021
0421432+193413	IRAS 04187+1927	4.274 ± 0.041	2.494 ± 0.027		10.185 ± 0.022	8.725 ± 0.024	8.021 ± 0.018
0421556+275506	DE Tau	5.382 ± 0.052	3.455 ± 0.017	12.773 ± 0.165	9.180 ± 0.022	8.273 ± 0.018	7.799 ± 0.018
0421574+282635	RY Tau	1.652 ± 0.017	-0.273 ± 0.019	10.064 ± 0.037	7.155 ± 0.019	6.128 ± 0.061	5.395 ± 0.023
0421588+281806	HD 283572	6.805 ± 0.274		8.847 ± 0.044	7.414 ± 0.029	7.008 ± 0.026	6.869 ± 0.023
0421594+193205	T Tau N (+Sab)	1.539 ± 0.019	-1.127 ± 0.029	9.747 ± 0.071	7.240 ± 0.023	6.237 ± 0.017	5.325 ± 0.017

Table 1. continued.

AKARI IRC name[a]	Identifier	AKARI *S9W*[b] [mag]	AKARI *L18W*[b] [mag]	UCAC[c] [mag]	2MASS *J* [mag]	2MASS *H* [mag]	2MASS K_S [mag]
0422007+265733	FS Tau B	5.327 ± 0.057	2.181 ± 0.316		15.082 ± 0.082	13.351 ± 0.053	11.753 ± 0.029
0422022+265729	FS Tau Aab	4.453 ± 0.044	1.877 ± 0.149	13.796 ± 0.900	10.705 ± 0.027	9.244 ± 0.026	8.178 ± 0.017
0422167+265456	CFHT-21	5.699 ± 0.065	3.793 ± 0.076	13.575 ± 1.000	11.577 ± 0.019	10.037 ± 0.022	9.011 ± 0.021
0422478+264552	IRAS 04196+2638	5.759 ± 0.099	4.330 ± 0.053		11.589 ± 0.022	10.147 ± 0.025	9.287 ± 0.019
0423077+280557	IRAS 04200+2759		3.998 ± 0.140		13.179 ± 0.021	11.600 ± 0.024	10.413 ± 0.018
0423354+250300	FU Tau	6.199 ± 0.024			10.781 ± 0.026	9.945 ± 0.027	9.324 ± 0.024
0423391+245613	FT Tau	6.030 ± 0.071	3.809 ± 0.128	14.791 ± 0.174	10.192 ± 0.026	9.123 ± 0.027	8.596 ± 0.021
0424570+271156	IP Tau	6.019 ± 0.039		12.046 ± 0.900	9.781 ± 0.021	8.893 ± 0.017	8.349 ± 0.018
0426535+260654	FV Tau AB	4.086 ± 0.027	2.116 ± 0.066	14.135 ± 0.900	9.917 ± 0.022	8.325 ± 0.024	7.442 ± 0.020
0426573+260629	KPNO-13	7.081 ± 0.157			11.281 ± 0.022	10.172 ± 0.022	9.580 ± 0.021
0427025+260531	DG Tau B	5.336 ± 0.027	1.838 ± 0.051		16.516 ± 0.141	13.525 ± 0.000	12.038 ± 0.000
0427028+254222	DF Tau AB	4.272 ± 0.054	2.644 ± 0.074	11.703 ± 0.075	8.171 ± 0.026	7.256 ± 0.023	6.734 ± 0.024
0427046+260616	DG Tau A	2.995 ± 0.017	0.369 ± 0.026		8.691 ± 0.018	7.722 ± 0.031	6.992 ± 0.020
0427572+261918	IRAS 04248+2612 AB	6.750 ± 0.105	3.307 ± 0.045		13.235 ± 0.053	11.795 ± 0.047	11.026 ± 0.040
0429208+274207	IRAS 04262+2735	4.379 ± 0.036	3.326 ± 0.065	15.095 ± 1.000	8.550 ± 0.026	7.461 ± 0.016	7.015 ± 0.017
0429217+270126	IRAS 04263+2654	6.545 ± 0.149	4.190 ± 0.107		10.801 ± 0.022	9.496 ± 0.022	8.725 ± 0.017
0429237+243301	GV Tau AB	1.579 ± 0.023	-1.163 ± 0.016		11.544 ± 0.025	9.587 ± 0.023	8.054 ± 0.024
0429300+243955	IRAS 04264+2433	5.838 ± 0.065	2.047 ± 0.056		13.728 ± 0.036	11.982 ± 0.031	11.135 ± 0.027
0429360+243554	J04293606+2435556	6.573 ± 0.041			10.780 ± 0.019	9.392 ± 0.020	8.659 ± 0.020
0429415+263258	DH Tau AB	6.332 ± 0.201	4.073 ± 0.070	13.462 ± 1.000	9.767 ± 0.021	8.824 ± 0.026	8.178 ± 0.026
0429515+260644	IQ Tau	5.274 ± 0.068	3.574 ± 0.070	13.890 ± 0.226	9.415 ± 0.020	8.417 ± 0.023	7.779 ± 0.023
0430039+181348	UX Tau C		3.478 ± 0.038		8.623 ± 0.023	7.960 ± 0.018	7.551 ± 0.021
0430296+242644	FX Tau AB	5.505 ± 0.079	3.556 ± 0.125	12.230 ± 0.900	9.388 ± 0.024	8.398 ± 0.018	7.924 ± 0.016
0430442+260124	DK Tau AB	4.191 ± 0.030	2.399 ± 0.033	11.986 ± 0.071	8.719 ± 0.030	7.758 ± 0.024	7.096 ± 0.016
0430502+230008	IRAS 04278+2253	2.103 ± 0.019	0.198 ± 0.020	15.580 ± 0.364	8.778 ± 0.024	7.040 ± 0.015	5.855 ± 0.023
0430513+244221	ZZ Tau AB	6.831 ± 0.153		14.544 ± 1.000	9.495 ± 0.021	8.695 ± 0.029	8.441 ± 0.021
0430517+244147	ZZ Tau IRS	5.883 ± 0.069	2.903 ± 0.103		12.842 ± 0.023	11.435 ± 0.026	10.314 ± 0.023
0431341+180804	L1551 IRS5	2.794 ± 0.026			13.708 ± 0.060	11.505 ± 0.052	9.822 ± 0.035
0431361+181344	LkHa 358	5.437 ± 0.170			12.792 ± 0.032	10.922 ± 0.029	9.687 ± 0.021
0431384+181357	HL Tau	2.697 ± 0.030	-0.231 ± 0.058		10.624 ± 0.042	9.171 ± 0.046	7.410 ± 0.017
0431400+181357	XZ Tau AB	3.421 ± 0.079		13.678 ± 0.124	9.385 ± 0.027	8.148 ± 0.040	7.291 ± 0.024
0431505+242418	HK Tau AB	6.255 ± 0.019	3.428 ± 0.060	13.588 ± 0.900	10.451 ± 0.022	9.253 ± 0.022	8.593 ± 0.018
0431578+182137	V710 Tau AB	5.999 ± 0.282	4.013 ± 0.194	13.560 ± 1.000	9.279 ± 0.032	9.107 ± 0.030	8.654 ± 0.025
0432154+242859	Haro 6-13	4.501 ± 0.093	1.604 ± 0.061		11.237 ± 0.024	9.319 ± 0.020	8.101 ± 0.029
0432303+173139	GG Tau Aa	4.540 ± 0.111	2.599 ± 0.051	11.785 ± 0.019	8.674 ± 0.035	7.815 ± 0.026	7.364 ± 0.018
0432317+242002	FZ Tau	4.138 ± 0.119	2.417 ± 0.030	13.352 ± 0.900	9.895 ± 0.022	8.400 ± 0.029	7.347 ± 0.017
0432320+225726	IRAS 04295+2251		1.947 ± 0.068		14.889 ± 0.044	11.982 ± 0.036	10.141 ± 0.024
0432429+255231	UZ Tau Aab	4.085 ± 0.022	2.247 ± 0.022	10.985 ± 1.000	9.136 ± 0.000	8.117 ± 0.000	7.354 ± 0.033
0432491+225303	JH112	5.189 ± 0.019	3.086 ± 0.047	15.093 ± 0.172	10.238 ± 0.027	8.995 ± 0.032	8.169 ± 0.023
0433062+240933	GH Tau AB	5.483 ± 0.137	3.804 ± 0.159	12.584 ± 1.000	9.109 ± 0.021	8.234 ± 0.027	7.794 ± 0.021
0433066+240955	V807 Tau AB	5.198 ± 0.089	3.638 ± 0.073	11.197 ± 1.000	8.146 ± 0.023	7.357 ± 0.026	6.960 ± 0.016
0433190+224633	IRAS 04303+2240	3.695 ± 0.060	2.000 ± 0.035	17.446 ± 1.000	11.103 ± 0.020	9.209 ± 0.023	7.673 ± 0.027
0433329+180059	HD 28867	5.092 ± 0.096	3.885 ± 0.256	6.466 ± 0.093	5.915 ± 0.035	5.813 ± 0.029	5.786 ± 0.023
0433341+242114	GI Tau	3.998 ± 0.127	1.848 ± 0.043	13.488 ± 1.000	9.341 ± 0.020	8.418 ± 0.021	7.888 ± 0.023

Table 1. continued.

AKARI IRC name[a]	Identifier	AKARI *S9W*[b] [mag]	AKARI *L18W*[b] [mag]	UCAC[c] [mag]	2MASS *J* [mag]	2MASS *H* [mag]	2MASS K_S [mag]
0433346+242106	GK Tau	4.478 ± 0.089		12.044 ± 1.000	9.053 ± 0.027	8.108 ± 0.026	7.468 ± 0.021
0433367+260949	IS Tau AB	5.662 ± 0.056	3.970 ± 0.047		10.323 ± 0.021	9.293 ± 0.023	8.642 ± 0.018
0433390+252038	DL Tau	4.749 ± 0.091	2.673 ± 0.041	12.894 ± 0.112	9.630 ± 0.021	8.679 ± 0.027	7.960 ± 0.021
0433394+175152	HN Tau AB	4.538 ± 0.061	2.574 ± 0.046	13.290 ± 0.124	10.699 ± 0.026	9.471 ± 0.027	8.384 ± 0.021
0433446+261500	SST Tau 043344.6+261500	6.791 ± 0.116			11.639 ± 0.021	10.385 ± 0.022	9.744 ± 0.018
0433519+225030	CI Tau	5.034 ± 0.093	2.926 ± 0.024	13.157 ± 0.142	9.480 ± 0.020	8.431 ± 0.040	7.793 ± 0.020
0433546+261326	IT Tau AB	5.661 ± 0.091	3.911 ± 0.169	13.073 ± 0.900	9.866 ± 0.025	8.591 ± 0.036	7.860 ± 0.026
0434554+242852	AA Tau	5.439 ± 0.017	3.556 ± 0.025	12.373 ± 1.000	9.433 ± 0.024	8.546 ± 0.023	8.047 ± 0.024
0435273+241458	DN Tau	5.816 ± 0.063	3.647 ± 0.120	12.171 ± 0.087	9.139 ± 0.021	8.342 ± 0.027	8.015 ± 0.021
0435410+241108	CoKu Tau/3 AB	5.129 ± 0.072	3.664 ± 0.043		10.731 ± 0.026	9.197 ± 0.026	8.411 ± 0.024
0435473+225021	HQ Tau	4.208 ± 0.064		11.788 ± 0.144	8.655 ± 0.024	7.731 ± 0.016	7.135 ± 0.021
0435528+225422	HP Tau AB	4.383 ± 0.090	2.049 ± 0.031	14.090 ± 0.201	9.549 ± 0.022	8.469 ± 0.065	7.625 ± 0.024
0435568+225436	Haro 6-28 AB	6.885 ± 0.072		15.986 ± 1.000	11.142 ± 0.022	10.055 ± 0.022	9.531 ± 0.020
0437514+262358	HT Tau	6.423 ± 0.143		12.859 ± 1.000	8.635 ± 0.026	7.542 ± 0.018	7.129 ± 0.018
0438285+261048	DO Tau	3.965 ± 0.050	1.706 ± 0.062		9.470 ± 0.022	8.243 ± 0.033	7.303 ± 0.017
0438352+261038	HV Tau AB+C	7.162 ± 0.264			9.227 ± 0.023	8.284 ± 0.026	7.906 ± 0.024
0439174+224753	VY Tau A	6.376 ± 0.123		13.385 ± 0.081	9.970 ± 0.023	9.260 ± 0.021	8.958 ± 0.020
0439178+222103	LkCa 15	5.834 ± 0.109	4.117 ± 0.123	11.875 ± 0.128	9.424 ± 0.020	8.600 ± 0.018	8.163 ± 0.018
0439208+254501	GN Tau B	5.405 ± 0.149	3.515 ± 0.027	15.465 ± 0.900	10.196 ± 0.025	8.893 ± 0.026	8.060 ± 0.027
0439557+254501	IC2087 IRS	2.631 ± 0.077	0.839 ± 0.084		10.668 ± 0.023	8.052 ± 0.018	6.275 ± 0.018
0440080+260524	IRAS 04370+2559	5.278 ± 0.237	3.081 ± 0.361		12.406 ± 0.023	10.248 ± 0.029	8.869 ± 0.018
0441168+283959	CoKu Tau/4		3.532 ± 0.069	14.771 ± 0.135	10.163 ± 0.030	9.077 ± 0.024	8.656 ± 0.019
0441387+255624	IRAS 04385+2550	5.488 ± 0.046	2.745 ± 0.039		11.849 ± 0.023	10.123 ± 0.022	9.200 ± 0.018
0442077+252310	V955 Tau Ab	4.937 ± 0.041	3.270 ± 0.018		9.811 ± 0.022	8.601 ± 0.021	7.942 ± 0.016
0442210+252033	CIDA-7	7.088 ± 0.058			11.397 ± 0.023	10.575 ± 0.022	10.169 ± 0.018
0442376+251537	DP Tau	4.898 ± 0.058	2.641 ± 0.056	15.058 ± 0.900	10.995 ± 0.021	9.689 ± 0.017	8.760 ± 0.016
0446530+165959	DQ Tau	5.093 ± 0.031	2.820 ± 0.065	13.253 ± 0.180	9.511 ± 0.021	8.544 ± 0.020	7.981 ± 0.021
0446590+170238	Haro 6-37 A	4.597 ± 0.019	2.747 ± 0.064	13.655 ± 1.000	9.239 ± 0.028	7.991 ± 0.021	7.310 ± 0.024
0447062+165842	DR Tau	3.552 ± 0.020		12.187 ± 0.410	8.845 ± 0.024	7.799 ± 0.053	6.874 ± 0.017
0447485+292511	DS Tau	5.621 ± 0.085	4.245 ± 0.050	12.490 ± 0.165	9.465 ± 0.018	8.597 ± 0.033	8.036 ± 0.029
0451473+304712	UY Aur A	3.479 ± 0.018	1.152 ± 0.016	12.151 ± 0.137	9.134 ± 0.020	7.987 ± 0.016	7.239 ± 0.018
0452066+304717	IRAS 04489+3042	5.826 ± 0.067	3.246 ± 0.041		14.426 ± 0.030	12.021 ± 0.021	10.383 ± 0.018
0452096+303744	Haro 6-39	7.023 ± 0.181	4.539 ± 0.110		13.254 ± 0.021	12.117 ± 0.018	11.187 ± 0.018
0455095+182629	HD 31281	7.221 ± 0.257		9.105 ± 0.088	7.974 ± 0.027	7.681 ± 0.017	7.609 ± 0.029
0455110+302159	GM Aur	6.793 ± 0.063	3.338 ± 0.131		9.341 ± 0.018	8.603 ± 0.024	8.283 ± 0.017
0455458+303303	AB Aur	1.185 ± 0.016	-0.841 ± 0.026	7.543 ± 0.157	5.936 ± 0.018	5.062 ± 0.020	4.230 ± 0.016
0455560+303622	XEST26-062	6.521 ± 0.154	4.230 ± 0.089	15.339 ± 0.054	10.471 ± 0.021	9.660 ± 0.018	9.267 ± 0.019
0455593+303401	SU Aur	3.445 ± 0.016	0.671 ± 0.017	9.057 ± 0.027	7.199 ± 0.020	6.558 ± 0.020	5.990 ± 0.023
0457065+314250	RXJ0457.0+3142	6.108 ± 0.055		10.002 ± 0.081	7.486 ± 0.021	6.756 ± 0.018	6.538 ± 0.023
0503066+252319	V836 Tau	6.378 ± 0.297		13.817 ± 0.203	9.913 ± 0.023	9.077 ± 0.029	8.595 ± 0.019
0505228+253131	CIDA-9	6.337 ± 0.062	4.870 ± 0.021	16.870 ± 1.000	12.808 ± 0.035	11.913 ± 0.043	11.161 ± 0.029
0507495+302404	RW Aur A	3.453 ± 0.047	1.620 ± 0.036	9.878 ± 0.076	8.378 ± 0.024	7.621 ± 0.038	7.020 ± 0.018
0529083+115212	1RXS J052908.4+115207	2.605 ± 0.017	0.168 ± 0.024	9.443 ± 0.049	7.698 ± 0.030	7.103 ± 0.029	6.590 ± 0.029
0529406+291110	1RXS J052940.9+291058	5.039 ± 0.037		6.260 ± 0.265	5.314 ± 0.020	5.208 ± 0.252	5.028 ± 0.021

Table 1. continued.

AKARI IRC name[a]	Identifier	AKARI *S9W*[b] [mag]	AKARI *L18W*[b] [mag]	UCAC[c] [mag]	2MASS J [mag]	2MASS H [mag]	2MASS K_S [mag]
0536516+232605	1RXS J053652.7+232600	6.789 ± 0.209		8.591 ± 0.032	7.328 ± 0.029	7.022 ± 0.021	6.913 ± 0.020
0537184+133452	1RXS J053718.4+133453	7.056 ± 0.016		9.377 ± 0.030	8.105 ± 0.024	7.720 ± 0.020	7.593 ± 0.020

[a] Source name from its J2000 coordinates, following the IAU Recommendations for Nomenclature (2006). The format is HHMMSSS±DDMMSS. The source must be referred to in the literatures by its full name: AKARI-IRC-V1 J0123456+765432, where V1 refers to the version code.

[b] The zero-magnitude flux densities for the AKARI bands are 56.262 Jy and 12.001 Jy for the *S9W* and *L18W* bands, respectively. The AKARI measurements are based on the V-1 version of the IRC-PSC.

[c] The UCAC magnitude is derived from aperture photometry.

[d] 1RXS J032409.7+123745 seems to be a giant star.

Table 2. AKARI, UCAC, and 2MASS photometric data for the new TTS candidates.

AKARI IRC name[a]	AKARI *S9W*[b] [mag]	AKARI *L18W*[b] [mag]	UCAC[c] [mag]	2MASS *J* [mag]	2MASS *H* [mag]	2MASS K_S [mag]
0322025+305129	5.761 ± 0.077		10.027 ± 0.022	8.227 ± 0.018	7.850 ± 0.020	7.663 ± 0.020
0325067+310652	6.473 ± 0.111	3.546 ± 0.125	12.613 ± 0.060	9.684 ± 0.022	8.930 ± 0.021	8.438 ± 0.018
0325125+305922	6.589 ± 0.154		13.913 ± 0.086	10.243 ± 0.021	9.414 ± 0.021	9.067 ± 0.020
0337114+330303	6.441 ± 0.109	4.488 ± 0.108	13.630 ± 0.212	10.631 ± 0.023	9.746 ± 0.021	9.223 ± 0.018
0349290+345800	6.227 ± 0.097	4.400 ± 0.242	13.742 ± 0.114	10.933 ± 0.022	10.082 ± 0.016	9.504 ± 0.018
0413573+291820	5.648 ± 0.044	3.559 ± 0.152	16.015 ± 0.316	11.310 ± 0.023	10.157 ± 0.024	9.356 ± 0.024
0414187+115812	6.330 ± 0.180	4.291 ± 0.194	12.456 ± 0.104	10.128 ± 0.022	9.554 ± 0.022	9.205 ± 0.018
0421156+100722	6.732 ± 0.101		11.685 ± 0.064	9.107 ± 0.020	8.409 ± 0.047	8.168 ± 0.026
0425412+353718		3.786 ± 0.254	13.669 ± 0.112	11.172 ± 0.021	10.545 ± 0.022	10.069 ± 0.017
0430375+355031	6.264 ± 0.178	3.827 ± 0.241	16.213 ± 0.372	10.027 ± 0.021	8.874 ± 0.018	8.052 ± 0.019
0435254+341901	6.208 ± 0.063	4.073 ± 0.163	15.805 ± 0.339	11.566 ± 0.022	10.339 ± 0.028	9.503 ± 0.021
0450190+092328	4.337 ± 0.033	2.876 ± 0.032	12.323 ± 0.900	7.118 ± 0.024	6.253 ± 0.024	5.970 ± 0.024
0502405+192237	4.908 ± 0.029	2.959 ± 0.086	13.449 ± 0.224	8.119 ± 0.030	7.013 ± 0.017	6.576 ± 0.017
0511021+295926	6.684 ± 0.138	4.367 ± 0.045	13.091 ± 0.057	10.063 ± 0.035	9.320 ± 0.041	9.022 ± 0.032
0512342+255847	6.145 ± 0.206		13.313 ± 0.129	10.462 ± 0.022	9.706 ± 0.022	9.411 ± 0.018
0516039+061852	6.807 ± 0.080		14.728 ± 0.112	11.286 ± 0.022	10.463 ± 0.027	9.978 ± 0.018
0517259+070022	5.130 ± 0.045	3.156 ± 0.036	11.000 ± 0.026	9.469 ± 0.039	8.941 ± 0.055	8.424 ± 0.034
0519413+053842	5.107 ± 0.031	2.086 ± 0.023	8.977 ± 0.018	8.041 ± 0.023	7.706 ± 0.023	7.482 ± 0.024
0522456+225444		2.467 ± 0.063	13.459 ± 0.114	9.641 ± 0.022	8.693 ± 0.026	8.391 ± 0.017
0525113+191547	6.331 ± 0.101		14.599 ± 0.143	11.309 ± 0.021	10.416 ± 0.026	9.918 ± 0.018
0525519+345228	3.100 ± 0.077	1.520 ± 0.122	14.503 ± 0.155	9.621 ± 0.022	8.499 ± 0.020	7.745 ± 0.023
0527433+031309	7.023 ± 0.067	4.304 ± 0.066	12.163 ± 0.096	10.378 ± 0.023	9.849 ± 0.024	9.708 ± 0.025
0527493+064638	3.568 ± 0.119		14.185 ± 0.231	8.312 ± 0.026	7.290 ± 0.057	6.462 ± 0.021
0529091+235902	5.960 ± 0.116	4.267 ± 0.107	14.452 ± 0.120	11.259 ± 0.024	10.368 ± 0.020	9.784 ± 0.018
0529592+121947		4.078 ± 0.188	12.949 ± 0.119	10.593 ± 0.020	9.836 ± 0.021	9.391 ± 0.018
0536009+113339	6.718 ± 0.072		11.543 ± 0.091	9.863 ± 0.026	9.263 ± 0.022	8.980 ± 0.023
0537531+372456	5.511 ± 0.070	3.949 ± 0.035	15.577 ± 0.228	10.543 ± 0.022	9.827 ± 0.026	9.313 ± 0.024
0539023+085612	6.947 ± 0.039		11.170 ± 0.037	9.744 ± 0.024	9.157 ± 0.024	8.736 ± 0.021

[a] Source name from its J2000 coordinates, following the IAU Recommendations for Nomenclature (2006). The format is HHMMSSS±DDMMSS. The source must be referred to in the literatures by its full name: AKARI-IRC-V1 J0123456+765432, where V1 refers to the version code.

[b] The zero-magnitude flux densities for the AKARI bands are 56.262 Jy and 12.001 Jy for the *S9W* and *L18W* bands, respectively. The AKARI measurements are based on the V-1 version of the IRC-PSC.

[c] The UCAC magnitude is derived from aperture photometry.